

## LIST OF CONTRIBUTORS

- Masazumi Harada* Department of Social Welfare Studies,  
Kumamoto Gakuen University, Japan
- Katsuhiko Keida* Associate Professor, Faculty of Letters,  
Kumamoto University, Japan
- Toshinori Kitamura  
and Fusako Kitamura* Department of Clinical Behavioural Sciences,  
Kumamoto University, Japan
- Ichiro Matsuda* Professor of Medical Genetics and Pediatrics,  
Health Science University of Hokkaido, The  
Research Institute of Personalized Health  
Science, Japan
- Toshiko Morita* School of Health Sciences, Course of Nursing,  
Kumamoto University, Japan
- Tadashi Saga* Professor, Center for Lifelong Learning,  
Kumamoto University, Japan
- Hideyuki Saya* Professor, Graduate School of Medical  
Sciences, Kumamoto University,  
Japan
- Eri Shinoki* Assistant Professor of Maternal and Child  
Nursing, Health Science University of  
Hokkaido, The Research Institute of  
Personalized Health Science, Japan
- Susumu Nakayama* Professor, Faculty of Design and  
Architecture, Aichi Sangyo University,  
Japan
- Hiroaki Taguchi* Professor, Faculty of Letters, Kumamoto  
University, Japan
- Takao Takahashi* Professor, Faculty of Letters, Kumamoto  
University, Japan

*Tomohiro Tanaka*

Associate Professor, Faculty of Letters,  
Kumamoto University, Japan

*Hideyuki Yahata*

Associate Professor, Faculty of Education,  
Kumamoto University, Japan



