

How to select a model if we know probabilities with interval uncertainty?

Probabilities
with interval
uncertainty

Vladik Kreinovich

Department of Computer Science, University of Texas at El Paso, El Paso,
Texas, USA

Received 22 August 2023
Revised 15 September 2023
Accepted 15 September 2023

Abstract

Purpose – When the probability of each model is known, a natural idea is to select the most probable model. However, in many practical situations, the exact values of these probabilities are not known; only the intervals that contain these values are known. In such situations, a natural idea is to select some probabilities from these intervals and to select a model with the largest selected probabilities. The purpose of this study is to decide how to most adequately select these probabilities.

Design/methodology/approach – It is desirable to have a probability-selection method that preserves independence. If, according to the probability intervals, the two events were independent, then the selection of probabilities within the intervals should preserve this independence.

Findings – The paper describes all techniques for decision making under interval uncertainty about probabilities that are consistent with independence. It is proved that these techniques form a 1-parametric family, a family that has already been successfully used in such decision problems.

Originality/value – This study provides a theoretical explanation of an empirically successful technique for decision-making under interval uncertainty about probabilities. This explanation is based on the natural idea that the method for selecting probabilities from the corresponding intervals should preserve independence.

Keywords Decision making under uncertainty, Interval uncertainty about probabilities, Independent events, Maximum likelihood approach

Paper type Research paper

1. Formulation of the problem

1.1 Need for indirect measurements and data processing

In many practical situations, we are interested in the quantity y that is difficult – or even impossible to measure directly. For example, we may be interested in tomorrow's temperature or in next year's gross domestic product (GDP). Since we cannot measure the quantity y directly, we need to measure it *indirectly*, i.e.:

- (1) find easier-to-measure quantities x_1, \dots, x_n which are related to y by a known dependence $y = f(x_1, \dots, x_n)$,

© Vladik Kreinovich. Published in *Asian Journal of Economics and Banking*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

The author is thankful to Thierry Denœux, Hung T. Nguyen, and Nguyen Ngoc Thach, for their encouragement and advice.

This work was funded by the National Science Foundation (No. 1623190) (A Model of Change for Preparing a New Generation for Professional Practice in Computer Science), HRD-1834620 and HRD-2034030 (CAHSI Includes), EAR-2225395 (Center for Collective Impact in Earthquake Science C-CIES) and the AT&T Fellowship in Information Technology. It was also supported by the program for the development of Scientific-Educational Mathematical Center of Volga Federal District (No. 075-02-2020-1478) and the Hungarian National Research, Development and Innovation Office (NRDI).



-
- (2) measure the values of these quantities, resulting in measurement results $\tilde{x}_1, \dots, \tilde{x}_n$, and
 - (3) compute the desired estimate $\tilde{y} = f(\tilde{x}_1, \dots, \tilde{x}_n)$ by applying the algorithm f to the results \tilde{x}_i of measuring x_i .

Computing \tilde{y} is an important particular case of *data processing*.

1.2 Need to find a model

In many practical situations, we know the function $f(x_1, \dots, x_n)$. For example, in celestial mechanics, we know how the future location y of a celestial body depends on the current location and velocity of this and other bodies. However, in many other practical situations, we do not know this dependence. In such cases, we need to determine this dependence from the experiments and/or observations. Specifically,

- (1) In several (K) cases, we know both the values $x_i^{(k)}$ of the inputs x_i and the value $y^{(k)}$ of the desired quantity y , and
- (2) We need to find the dependence $f(x_1, \dots, x_n)$ that is consistent with all these observations, i.e. for which, for all k from 1 to K , we have

$$y^{(k)} = f(x_1^{(k)}, \dots, x_n^{(k)}). \quad (1)$$

1.2.1 Terminological comment.

- (1) The resulting function $y = f(x_1, \dots, x_n)$ serves as a *model* of the corresponding situation.
- (2) In statistics, the problem of finding a model is known as *regression*.
- (3) In computer science, the same problem – when solved by an algorithm – is known as *machine learning*.
- (4) In this paper, we use the word “model” in the general scientific sense – as a description of a real-life process, i.e. in this case, as a function $f(x_1, \dots, x_n)$ that estimates the desired quantity y . To avoid possible confusion, it should be mentioned that in statistics, sometimes, a “model” means a *family* of such functions – e.g. all linear functions or all linear functions that depend only on the first k variables x_1, \dots, x_k .

1.3 Need to select a model

- (1) To describe a general function $f(x_1, \dots, x_n)$, we need to describe infinitely many parameters – e.g. the values of the function at all the tuples (x_1, \dots, x_n) for which all the values x_i are rational.
- (2) Our only requirement on possible functions is to satisfy *K equations (1)*.

Here, the number of parameters much larger than the number of equations. Thus, there are, in general, many different functions that fit all the observations.

We therefore need to select one of these functions, i.e. we need to select a model.

1.4 How a model is selected now: case when we know probabilities

In some cases, we know the probabilities p_i of different models. In this case, a reasonable idea is to select the *most probable* model, i.e. the model whose probability is the largest: $p_i = \max_j p_j$.

Such a selection is, e.g. one of the main ideas behind the maximum likelihood approach to model selection; see, e.g. ([Sheskin, 2011](#)). In this method, usually, we maximize the probability p by solving the equivalent problem of minimizing the quantity

$$L \stackrel{\text{def}}{=} -\ln(p).$$

Probabilities
with interval
uncertainty

1.4.1 Comment. It should be mentioned that, strictly speaking, likelihood is not the probability of a model; it is the probability of the data according to this model. To come up with the probability of the model, we need to use Bayesian approach. In this approach, if we assume that *a priori* all models are equally probable – i.e. that prior distribution is uniform – then likelihood becomes proportional to the probability of the model, so that maximizing likelihood is equivalent to maximizing the model's probability.

1.5 What if we only have partial information about probabilities: description of the situation

Often, we only have partial information about the probabilities. For example, instead of the exact values p_i of each probability, we only know the lower bound \underline{p}_i and the upper bound \bar{p}_i : $\underline{p}_i \leq p_i \leq \bar{p}_i$.

In this case, the only information that we have about the probability p_i is that this probability is contained on the interval $[\underline{p}_i, \bar{p}_i]$. Thus, this situation is known as *interval uncertainty*.

1.6 How to make a decision under such interval uncertainty: a natural idea

In situations with interval uncertainty, it is desirable to apply well-traditional probability-based decision making technique. To do this, we need to select, within each of the intervals $[\underline{p}_i, \bar{p}_i]$, one of the values p_i , and then select the model with the largest value of this selected probability p_i .

1.7 Resulting challenge

How do we select a value p_i in each interval? There are many different ways to select, which one should we choose?

1.8 What we do in this paper

In this paper, we show that a natural condition on the selection of the probability values from the corresponding intervals uniquely determine a 1-parametric family of such selections – the only selections that satisfy this natural condition.

2. Main result

2.1 Natural condition: informal description

We want to find a mapping that assigns, to each interval of probability values, a number from this interval. It is desirable to select this mapping so that it preserves important properties of the situation.

In probabilistic techniques, one of the most important notions is the notion of independence. It is therefore reasonable to require that the desired intervals-to-numbers mapping satisfy the following condition: If the two events were independent, then this mapping should preserve this independence.

2.2 Let us formalize this natural condition

If two events with probabilities p_1 and p_2 are independent, then the probability of them occurring at same time is equal to the product $p_1 \cdot p_2$ of the corresponding probabilities. If for each of these events, we only know the interval $[\underline{p}_i, \bar{p}_i]$ of possible values of its probability, then possible values of the probability that both events occur is equal to the set of possible values

$$\left\{ p_1 \cdot p_2 : p_1 \in [\underline{p}_1, \bar{p}_1] \text{ and } p_2 \in [\underline{p}_2, \bar{p}_2] \right\}$$

one can easily check that this set is equal to the interval $[\underline{p}_1 \cdot \underline{p}_2, \bar{p}_1 \cdot \bar{p}_2]$; see, e.g. (Jaulin *et al.*, 2012), (Kubica, 2019; Mayer, 2017; Moore *et al.*, 2009). Indeed, for non-negative values p_i , the product function $p_1, p_2 \mapsto p_1 \cdot p_2$ is (nonstrictly) increasing with respect to each of its variables. Thus:

- (1) The smallest possible value of this function when $p_i \in [\underline{p}_i, \bar{p}_i]$ is attained when both inputs are the smallest possible, i.e. when $p_i = \underline{p}_i$ for both i , and
- (2) The largest possible value of this function when $p_i \in [\underline{p}_i, \bar{p}_i]$ is attained when both inputs are the largest possible, i.e. when $p_i = \bar{p}_i$ for both i .

Thus, we arrive at the following [definition](#).

Definition. We say that a mapping f that maps each subinterval $[\underline{p}, \bar{p}]$ of the interval $[0, 1]$ into a number $f(\underline{p}, \bar{p})$ from this interval is *natural* if it satisfies the following condition: for all values $\underline{p}_1 \leq \bar{p}_1$ and $\underline{p}_2 \leq \bar{p}_2$, we have

$$f(\underline{p}_1 \cdot \underline{p}_2, \bar{p}_1 \cdot \bar{p}_2) = f(\underline{p}_1, \bar{p}_1) \cdot f(\underline{p}_2, \bar{p}_2)$$

Proposition. A mapping is natural if and only if, for some $\alpha \in [0, 1]$, it has the form

$$f(\underline{p}, \bar{p}) = \underline{p}^\alpha \cdot \bar{p}^{1-\alpha}.$$

Discussion. The function $L = -\ln(p)$ is decreasing with respect to p . Thus, when $p \in [\underline{p}, \bar{p}]$, then:

- (1) The smallest value \underline{L} of $L = -\ln(p)$ is attained when p is the largest, i.e. when $p = \bar{p}$:

$$\underline{L} = -\ln(\bar{p});$$

- (2) The largest value \bar{L} of L is attained when p is the smallest, i.e. when $p = \underline{p}$:

$$\bar{L} = -\ln(\underline{p}).$$

For the values $L = -\ln(p)$, $\underline{L} = -\ln(\bar{p})$, and $\bar{L} = -\ln(\underline{p})$, the above formula takes the form $L = \alpha \cdot \bar{L} + (1 - \alpha) \cdot \underline{L}$. Interestingly, this is exactly Hurwicz optimism-pessimism criterion that is used for decision making under interval uncertainty; see, e.g. (Hurwicz, 1951; Kreinovich, 2014; Luce and Raiffa, 1989).

This model selection has been successfully used; see, e.g. [\(Denœux, 2023\)](#).

Probabilities
with interval
uncertainty

Proof of the Proposition.

1. It is easy to prove that the above formula leads to a natural mapping. So, to complete the proof, it is sufficient to prove that every natural mapping has this form.

Let $f(\underline{p}, \bar{p})$ be a natural mapping. Let us prove that it has the desired form.

2. For each p , by [definition](#) of a natural mapping, the value $f(p, p)$ belongs to the interval $[p, \bar{p}]$ and is, thus, equal to p . In particular, for $p = 0$, we get

$$f(0, 0) = 0.$$

3. Let us first take $\underline{p}_1 = \underline{p}_2 = 0$ and $\bar{p}_1 = \bar{p}_2 = 1$. In this case, the naturalness condition implies that $f(0, 1) \cdot f(0, 1) = f(0, 1)$. Thus, either $f(0, 1) = 1$ or $f(0, 1) = 0$. Let us consider these two possible cases one by one.

4. Let us first consider the case when $f(0, 1) = 1$.

4.1. In this case, for every $a \in [0, 1]$, for $\underline{p}_1 = 0$, $\bar{p}_1 = 1$, $\underline{p}_2 = 1$, and $\bar{p}_2 = 1$, we get $f(0, 1) \cdot f(a, 1) = f(0, 1)$. Since $f(0, 1) = 1$, this means that $f(a, 1) = 1$ for all a .

4.2. Now, for all possible $\underline{p} \leq \bar{p}$ for which $\bar{p} > 0$, naturalness leads to

$$f(\underline{p}, \bar{p}) = f(\bar{p}, \bar{p}) \cdot f(\underline{p}/\bar{p}, 1)$$

As we have proven in Section 4.1 of this proof, the second factor $f(\underline{p}/\bar{p}, 1)$ is equal to 1. The first factor $f(\bar{p}, \bar{p})$ is, by Part 2 of this proof, equal to \bar{p} .

So, for all cases when $\bar{p} > 0$, we have $f(\underline{p}, \bar{p}) = \bar{p}$.

4.3. For $\bar{p} = 0$, the formula $f(\underline{p}, \bar{p}) = \bar{p}$ is also true – by Part 2 of this proof. Thus, this formula holds for all $\underline{p} \leq \bar{p}$. This corresponds to $\alpha = 0$.

5. Let us now consider the case when $f(0, 1) = 0$.

In this case, naturalness implies that, for all \bar{p} , we have

$$f(0, \bar{p}) = f(0, 1) \cdot f(0, \bar{p})$$

and hence $f(0, \bar{p}) = 0$. Let us now consider intervals for which $\underline{p} > 0$, and thus, for $F(A) \stackrel{\text{def}}{=} -\ln(f(\exp(-A), 1))$, we always have $F(A) \geq 0$. In particular, $F(1) \geq 0$.

5.1. Let us first consider the values $f(a, 1)$ corresponding to $a > 0$. When $a < b$, then we have $f(a, 1) = f(a/b, 1) \cdot f(b, 1)$. Since $f(a/b, 1)$ is a probability, it is smaller than or equal to 1, thus, $f(a, 1) \leq f(b, 1)$, i.e. $f(a, 1)$ is a nonstrictly increasing function of a .

5.2. Each value $a > 0$ can be represented as $\exp(-A)$ for $A = -\ln(a)$. By [definition](#) of the natural mapping, each such value $f(a, 1)$ for $a > 0$ is greater than or equal to $a > 0$ and thus, $f(a, 1) > 0$. So, we can take logarithm of these values as well. Let us denote $F(A) \stackrel{\text{def}}{=} -\ln(f(\exp(-A), 1))$. Probabilities $f(\exp(-A), 1)$ are smaller than or equal to 1, so

$$\ln(f(\exp(-A), 1)) \leq 1 = 0$$

and thus, for $F(A) \stackrel{\text{def}}{=} -\ln(f(\exp(-A), 1))$, we always have $F(A) \geq 0$. In particular, $F(1) \geq 0$.

5.3. Let us prove that $F(A)$ is a (nonstrictly) increasing function.

Indeed, if $A < B$, then $-A > -B$. Since $\exp(x)$ is an increasing function, we get $\exp(-A) > \exp(-B)$. Since $f(a, 1)$ is a nonstrictly increasing function of a , we conclude that $f(\exp(-A), 1) \geq f(\exp(-B), 1)$. Since $\ln(x)$ is an increasing function, we conclude that

$$\ln(f(\exp(-A), 1)) \geq \ln(f(\exp(-B), 1)).$$

Multiplying both sides by -1 , we get

$$-\ln(f(\exp(-A), 1)) \leq -\ln(f(\exp(-B), 1)),$$

i.e. $F(A) \leq F(B)$. The statement is proven.

5.4. For values $f(a, 1)$, naturalness implies that $f(a \cdot b, 1) = f(a, 1) \cdot f(b, 1)$. For $a = \exp(-A)$ and $b = \exp(-B)$, we have $a \cdot b = \exp(-(A + B))$, thus,

$$f(\exp(-(A + B)), 1) = f(\exp(-A), 1) \cdot f(\exp(-B), 1).$$

By taking negative logarithms of both sides, we get

$$F(A + B) = F(A) + F(B). \quad (2)$$

5.5. For every integer m , formula (2) implies that

$$\begin{aligned} F(m \cdot A) &= F(A + \cdots + A \text{ (m times)}) = \\ F(A) + \cdots + F(A) \text{ (m times)} &= m \cdot F(A). \end{aligned} \quad (3)$$

in particular, for $m = n$ and $A = 1/n$, we get $F(1) = n \cdot F(1/n)$, hence

$$F(1/n) = (1/n) \cdot F(1). \quad (4)$$

for a general m and $A = 1/n$, we get $F(m/n) = m \cdot F(1/n)$. Due to (4), we get $F(m/n) = (m/n) \cdot F(1)$, i.e. $F(r) = r \cdot F(1)$ for all rational number r .

5.6. For every real number x and for every positive integer n , we can take, as m_n , an integer part of $n \cdot x$, so that $m_n \leq n \cdot x < m_n + 1$. By dividing all parts of this inequality by n , we get $m_n/n \leq x < (m_n + 1)/n$. In the limit $n \rightarrow \infty$, we get $m_n/n \rightarrow x$ and $(m_n + 1)/n \rightarrow x$.

By Part 5.3 of this proof, the function $F(A)$ is nonstrictly increasing, thus $F(m_n/n) \leq F(x) \leq F((m_n + 1)/n)$. Due to Part 5.5, this means that

$$(m_n/n) \cdot F(1) \leq F(x) \leq ((m_n + 1)/n) \cdot F(1).$$

In the limit $n \rightarrow \infty$, we have

$$(m_n/n) \cdot F(1) \rightarrow x \cdot F(1) \text{ and } ((m_n + 1)/n) \cdot F(1) \rightarrow x \cdot F(1).$$

Thus, in the limit, we get $x \cdot F(1) \leq F(x) \leq x \cdot F(1)$, i.e. $F(x) = x \cdot F(1)$.

5.7. By definition, $F(A) = -\ln(f(\exp(-A), 1))$, thus,

$$f(\exp(-A), 1) = \exp(-F(A)) = \exp(-A \cdot F(1)).$$

substituting $A = -\ln(a)$ into this expression, we get

$$f(a, 1) = e^{\ln(a) \cdot F(1)} = (e^{\ln(a)})^{F(1)} = a^{F(1)}.$$

Probabilities
with interval
uncertainty

The condition that $f(a, 1) \geq a$ implies that $F(1) \leq 1$, thus $F(1) \in [0, 1]$.

5.8. For every pair $0 < \underline{p} \leq \bar{p}$, naturalness implies that

$$f(\underline{p}, \bar{p}) = f(\bar{p}, \bar{p}) \cdot f(\underline{p}/\bar{p})$$

by Part 2 of this proof, the first factor in this product is equal to \bar{p} . Due to Part 5.7, we get the expression for the second factor, thus we get

$$f(\underline{p}, \bar{p}) = \bar{p} \cdot (\underline{p}/\bar{p})^{F(1)} = \underline{p}^{F(1)} \cdot \bar{p}^{1-F(1)}.$$

This is exactly the desired formula, for $\alpha = F(1)$ – limited to the case when $\underline{p} > 0$. Then:

- (1) If $\alpha = 0$, we get the case considered in Part 4 of this proof.
- (2) For $\alpha > 0$ and $\underline{p} = 0$, we have $0^\alpha \cdot \bar{p}^{1-\alpha} = 0$, and $f(0, \bar{p}) = 0$ by Part 5, so the desired equality holds for all $\underline{p} \leq \bar{p}$.

The proposition is proven.

References

- Denceux, T. (2023), “Quantifying prediction uncertainty in regression using random fuzzy sets: the ENNreg model”, *IEEE Transactions on Fuzzy Systems*, Vol. 31 No. 10, pp. 3690-3699, to appear, doi: [10.1109/tfuzz.2023.3268200](https://doi.org/10.1109/tfuzz.2023.3268200).
- Hurwicz, L. (1951), “Optimality criteria for decision making under ignorance”, Cowles Commission Discussion Paper, Statistics No. 370.
- Jaulin, L., Kiefer, M., Didrit, O. and Walter, E. (2012), *Applied Interval Analysis, with Examples in Parameter and State Estimation, Robust Control, and Robotics*, Springer, London.
- Kreinovich, V. (2014), “Decision making under interval uncertainty (and beyond)”, in Guo, P. and Pedrycz, W. (Eds), *Human-Centric Decision-Making Models for Social Sciences*, Springer Verlag, pp. 163-193.
- Kubica, B.J. (2019), *Interval Methods for Solving Nonlinear Constraint Satisfaction, Optimization, and Similar Problems: From Inequalities Systems to Game Solutions*, Springer, Cham.
- Luce, R.D. and Raiffa, R. (1989), *Games and Decisions: Introduction and Critical Survey*, Dover, New York.
- Mayer, G. (2017), *Interval Analysis and Automatic Result Verification*, de Gruyter, Berlin.
- Moore, R.E., Kearfott, R.B. and Cloud, M.J. (2009), *Introduction to Interval Analysis*, SIAM, Philadelphia.
- Sheskin, D.J. (2011), *Handbook of Parametric and Nonparametric Statistical Procedures*, Chapman and Hall/CRC, Boca Raton, FL.

Corresponding author

Vladik Kreinovich can be contacted at: vladik@utep.edu

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgroupublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com