

Structure theorem for Jordan algebra bundles

Ranjitha Kumar

Department of Mathematics, REVA University, Bengaluru, India

256

Received 13 August 2022
Revised 5 December 2022
3 March 2023
13 May 2023
Accepted 13 May 2023

Abstract

Purpose – The aims of this paper is to prove that every semisimple Jordan algebra bundle is locally trivial and establish the decomposition theorem for locally trivial Jordan algebra bundles using the decomposition theorem of Lie algebra bundles.

Design/methodology/approach – Using the decomposition theorem of Lie algebra bundles, this paper proves the decomposition theorem for locally trivial Jordan algebra bundles.

Findings – Findings of this paper establish the decomposition theorem for locally trivial Jordan algebra bundles.

Originality/value – To the best of the author's knowledge, all the results are new and interesting to the field of Mathematics and Theoretical Physics community.

Keywords Jordan algebra bundle, Lie algebra bundle, Vector bundle

Paper type Research paper

1. Introduction

In modern mathematics, an important notion is that of non-associative algebra. In Ref. [1], we gave a relationship between two important classes of non-associative algebras, namely, Lie algebras (introduced in 1870 by the Norwegian mathematician Sophus Lie in his study of the groups of transformations) and Jordan algebras (introduced in 1932–1933 by the German physicist Pasqual Jordan (1902–1980) in his algebraic formulation of quantum mechanics [2–4]). These two algebras are interconnected, as was remarked for instance by Kevin McCrimmon [5, p. 622]:

We are saying that if you open up a Lie algebra and look inside, 9 times out of 10 there is a Jordan algebra (of pair) which makes it work.

Here, we recall some connections between Jordan algebra bundles and Lie algebra bundles [1, 6]. If ξ is a locally trivial Jordan algebra bundle in which each fibre ξ_x has a unit element then

$$K(\xi) = \bigcup_{x \in X} K(\xi_x) = \bigcup_{x \in X} \left(\text{Der} \xi_x \oplus L(\xi_x) \oplus \xi_x \oplus \overline{\xi_x} \right)$$

is a Lie algebra bundle, where $\text{Der} \xi_x$ is the vector space of all derivation defined on ξ_x , $L(\xi_x)$ is the vector space of all left translations of ξ_x and $\overline{\xi_x}$ an isomorphic copy of ξ_x .

JEL Classification — 17Bxx, 17Cxx, 55Rxx

© Ranjitha Kumar. Published in the *Arab Journal of Mathematical Sciences*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) license. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this license may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

The author would like to thank the referee for constructive remarks that improve the presentation of the paper and for spotting several errors in the previous version of the paper. Also, the author would like to thank REVA University for its continuous support and encouragement.



Lemma 1.1. [1] Let ξ be a locally trivial Jordan algebra bundle over X , with each fibre ξ_x having a unit element e_x , $x \in X$. Then Jordan algebra bundles

$$g(\xi) = \bigcup_{x \in X} g(\xi_x) = \bigcup_{x \in X} (Der \xi_x + L(\xi_x))$$

is a Lie algebra bundle.

Theorem 1.2. [1] If ξ is a locally trivial Jordan algebra bundle in which each fibre has a unit element, then $K(\xi) = g(\xi) \oplus \xi \oplus \bar{\xi}$, is a Lie algebra bundle, where $\bar{\xi}$ is an isomorphic copy of the Jordan algebra bundle ξ . Further, ξ can be imbedded in $K(\xi)$ such that the Jordan multiplication on ξ can be given in terms of the Lie multiplication on $K(\xi)$.

Given a locally trivial Jordan algebra bundle ξ in which each fibre ξ_x has a unit element, consider the Lie algebra bundle $K(\xi) = g(\xi) \oplus \xi \oplus \bar{\xi}$. Let $h(\xi) = \bigcup_{x \in X} h(\xi_x)$, where each $h(\xi_x) = L(\xi_x) \oplus [L(\xi_x), L(\xi_x)]$ is an ideal of $g(\xi_x) = Der \xi_x + L(\xi_x)$ [7]. Let $\phi : U \times J \rightarrow p^{-1}(U)$ be the local triviality of the Jordan algebra bundle ξ , where J is a Jordan algebra. Set $g(\phi) = End\phi|_{U \times g(J)}$, where the vector bundle morphism $End\phi : U \times EndJ \rightarrow \bigcup_{x \in U} End \xi_x$ is given by $(End\phi)(x, f) = \phi_x f \phi_x^{-1}$. Further, $h(\xi)$ is a vector bundle since $g(\phi)|_{U \times h(J)}$ maps $U \times h(J)$ onto $\bigcup_{x \in U} h(\xi_x)$. Hence $h(\xi)$ is an ideal subbundle of $g(\xi)$. We denote by $L(\xi)$, the ideal subbundle $L(\xi) = h(\xi) \oplus \xi \oplus \bar{\xi}$ of $K(\xi)$. \square

In this paper, we prove that any semisimple Jordan algebra bundle is locally trivial and we supply an example to show that the converse need not be true. Further, we prove that a semisimple Jordan algebra bundle can be written as the direct sum of simple ideal bundles.

1.1 Notations and terminology

All Jordan algebra bundles $\xi = (\xi, p, X, \theta)$ are over the arbitrary topological space X unless otherwise mentioned.

2. Preliminaries

Definition 2.1. A Jordan algebra bundle is a vector bundle $\xi = (\xi, p, X)$ together with a vector bundle morphism $\theta : \xi \otimes \xi \rightarrow \xi$ inducing a Jordan algebra structure on each fibre ξ_x , $x \in X$.

Definition 2.2. By a trivial Jordan algebra bundle, we mean a trivial vector bundle $(X \times J, p, X)$, where J is a Jordan algebra.

Definition 2.3. A morphism $f : \xi \rightarrow \zeta$ of Jordan bundles ξ and ζ is a morphism of the underlying vector bundles such that for every $x \in X$, $f_x : \xi_x \rightarrow \zeta_x$ is a Jordan algebra homomorphism. If f is bijective and f^{-1} is continuous, then f is called an isomorphism.

Definition 2.4. By a subalgebra (ideal) bundle of a Jordan algebra bundle $\xi = (\xi, p, X)$, we mean a vector sub-bundle $\xi' = (\xi', p, X)$ of ξ such that each fibre $(\xi')_x$ is a subalgebra (ideal) of ξ_x , $\forall x \in X$.

Definition 2.5. By a semisimple Jordan algebra bundle, we mean a Jordan algebra bundle in which each fibre is a semisimple Jordan algebra.

Definition 2.6. If ξ is a Jordan algebra bundle with a nontrivial multiplication $\theta : \xi \otimes \xi \rightarrow \xi$ inducing the Jordan algebra bundle structure and if ξ has no ideal bundles except itself and the zero bundle, then we call ξ a simple Jordan algebra bundle.

Definition 2.7. A Jordan algebra bundle ξ is said to be the direct sum of the ideal bundles $\xi_1, \xi_2, \dots, \xi_n$ provided, $\xi = \xi_1 \oplus \xi_2 \oplus \dots \oplus \xi_n$

3. Semisimple Jordan algebra bundles

Definition 3.1. A Jordan algebra bundle $\xi = (\xi, p, X)$ is locally trivial if, for every open set $U \subseteq X$, there exists a Jordan algebra J , together with a (vector bundle) trivialization $\varphi : \xi|_U \rightarrow U \times J$, which is fibrewise a homeomorphism of Jordan algebras.

Not every Jordan algebra bundle is locally trivial, as the following example shows.

Example 3.2. Let J be a nonzero real Jordan algebra and \mathbb{R} the field of real numbers. Then $\xi = (\mathbb{R} \times J, p, \mathbb{R})$ is a trivial vector bundle. Let $\theta : \xi \otimes \xi \rightarrow \xi$ be defined by, $\theta(t, (u, v)) = t(uv)$ for all $t \in \mathbb{R}, u, v \in J$. Then, θ is continuous and induces a Jordan algebra structure on each fibre $\xi_x = x \times J$. Hence, ξ is a Jordan algebra bundle. But ξ is not locally trivial because ξ_t for $t \neq 0$ is a nonzero algebra whereas ξ_0 is the zero algebra.

Remark 3.3. From Atiyah [8, p. 4], we reproduce the following things without any changes. Suppose that V and W are vector spaces and that $E = X \times V$ and $F = X \times W$ are the corresponding product bundles. Then $\phi : E \rightarrow F$ determines a map $\Phi : X \rightarrow \text{Hom}(V, W)$ by formula $\Phi(x)(v) = \phi(v)$. Moreover, if we give $\text{Hom}(V, W)$ its usual topology, then Φ is continuous; conversely, any such continuous map $\Phi : X \rightarrow \text{Hom}(V, W)$ determines a homomorphism $\phi : E \rightarrow F$.

Theorem 3.4. Every semisimple Jordan algebra bundle is locally trivial.

Proof. Let ξ be a semisimple Jordan algebra bundle. The local triviality of ξ as a vector bundle is given by the vector bundle isomorphism $\alpha : U \times V \rightarrow p^{-1}(U)$. Then $\theta : \xi \otimes \xi \rightarrow \xi$ induces the morphism $\hat{\theta} : U \times (V \otimes V) \rightarrow U \times V$ given by,

$$\hat{\theta}(x, (v, w)) = \alpha_x^{-1} \theta(\alpha_x(v), \alpha_x(w)). \tag{3.1}$$

Then, from above Remark (3.3), $x \mapsto \hat{\theta}_x$ defines a continuous mapping from U to $M \subseteq \text{Hom}(V \times V, V)$, the space of all Jordan multiplications defined on V endowed with the subspace topology. Since the Jordan algebra $J_0 = (V, \hat{\theta}_{x_0})$, $x_0 \in U$ is semisimple, J_0 is rigid by [9, Corollary 1.3]. That is the orbit

$$G(x_0) = \{g \cdot \theta_{x_0} \mid g \in \text{Aut}(V)\},$$

with respect to the Lie group $G = \text{Aut}(V)$ is open in M .

Let $U' = \{x \in U \mid \hat{\theta}_x \in G(x_0)\}$. Then, for each $x \in U'$, there exists a g_x in G such that

$$\hat{\theta}_x = g_x \cdot \hat{\theta}_{x_0} = g_x \hat{\theta}_{x_0} (g_x^{-1} \oplus g_x^{-1}). \tag{3.2}$$

Further, since G and $G(x_0)$ satisfy the hypothesis of Aren's theorem [10] G/G_0 is homeomorphic to $G(x_0)$, where G_0 is the stability subgroup corresponding to $\hat{\theta}_{x_0}$. Also $G \rightarrow G/G_0$ is a principal bundle [11, p. 33] together with a local cross section given by $gG_0 \rightarrow g$ [12, p. 126]. Hence, the map $x \mapsto g_x$ becomes the composition of the continuous maps

$$x \mapsto \hat{\theta}_x \mapsto g_x G_0 \mapsto g_x.$$

Therefore, we can define the vector bundle isomorphism $\phi : U \times J_0 \rightarrow p^{-1}(U)$ by $\phi(x, v) = \alpha_x(g_x(v))$. The map ϕ preserves the Jordan multiplication

$$\begin{aligned}
 \phi\left(x, \widehat{\theta}_{x_0}(a, b)\right) &= \alpha_x g_x \widehat{\theta}_{x_0}(a, b) \\
 &= \alpha_x \widehat{\theta}_x(g_x(a), g_x(b)) \quad \text{by (3.2)} \\
 &= \alpha_x \alpha_x^{-1} \theta_x(\alpha_x g_x(a), \alpha_x g_x(b)) \quad \text{by (3.1)} \\
 &= \theta_x(\phi(x, a), \phi(x, b)).
 \end{aligned}$$

Hence, ϕ gives the required local triviality of the Jordan algebra bundle ξ . \square

4. Decomposition theorem for Jordan algebra bundle

Lemma 4.1. *Let $(X \times V, q, X)$ be a trivial vector bundle and $(X \times J, p, X)$ a trivial semisimple Jordan bundle. Suppose $\phi : X \times V \rightarrow X \times J$ is a vector bundle monomorphism such that for each $x \in X$, $\phi(x, V)$ is an ideal in J . Then, there exists a finite open partition $\cup_i X_i = X$ such that $\phi_x(V) = \phi_y(V)$ for $x, y \in X_i$. In particular, if X is connected, for all $x, y \in X$, $\phi_x(V) = \phi_y(V)$.*

Proof. The map $\phi : X \times V \rightarrow X \times J$ being a vector bundle morphism, $x \mapsto \phi_x$ is a continuous map from X to $\text{Hom}(V, J)$, the vector space of all linear transformations from V to J . If \tilde{I} denotes the collection of all distinct ideals of J whose dimension is equal to that of V , then by the semisimplicity of J , \tilde{I} is a finite set [13, Corollary.4.6, p. 98]. Let $\tilde{I} = \{I_1, I_2, \dots, I_n\}$ and let $X_i = \{x \in X \mid \phi_x(V) = I_i\}$ for $i = 1, 2, \dots, n$. Let $x \in X$. Since $\phi_x(V)$ is an ideal in J , and $\dim \phi_x(V) = \dim V$, we have $\phi_x(V) = I_i$ for some i . Thus, $x \in X_i$, and consequently $X = \cup_i X_i$.

It is enough to prove that each X_i is open in X . Let \tilde{J}_i be the vector subspace $\tilde{J}_i = \{T \in \text{Hom}(V, J) \mid T(V) \subseteq I_i\}$. Then \tilde{J}_i is a closed subset of $\text{Hom}(V, J)$ being a linear subspace of the vector space $\text{Hom}(V, J)$. Since $x \mapsto \phi_x$ is continuous and \tilde{J}_i is closed in $\text{Hom}(V, J)$, $\tilde{X}_i = \{x \in X \mid \phi_x \in \tilde{J}_i\}$ is closed in X .

To prove: $\tilde{X}_i = X_i$

$x \in X_i \Rightarrow \phi_x(V) = I_i \Rightarrow \phi_x \in \tilde{J}_i \Rightarrow x \in \tilde{X}_i \Rightarrow X_i \subseteq \tilde{X}_i$. Now if $x \in \tilde{X}_i$, $\phi_x \in \tilde{J}_i$. Then, $\phi_x(V) \subseteq I_i$. But since $\dim \phi_x(V) = \dim I_i$, $\phi_x(V) = I_i$. So $x \in X_i$. Thus, $X_i = \tilde{X}_i$.

Consider $\tilde{X}_i \cap \tilde{X}_j$ and let $i \neq j$

$$x \in \tilde{X}_i \cap \tilde{X}_j \Rightarrow \phi_x \in \tilde{J}_i \text{ and } \phi_x \in \tilde{J}_j \Rightarrow \phi_x(V) \subseteq I_i \cap I_j = \emptyset$$

Therefore, $\tilde{X}_i \cap \tilde{X}_j = \emptyset$

Thus, \tilde{X}_i 's are all disjoint collection of closed sets. Hence, X_i is the complement of $\cup_{i \neq j} \tilde{X}_j$. So X_i is open in X . We here note that if X is connected, then for all $x, y \in X$, $\phi_x(V) = \phi_y(V)$. \square

Lemma 4.2. *Let ξ' be an ideal bundle of a semisimple Jordan algebra bundle ξ . Then, $h_1(\xi') = \cup_{x \in X} h_1(\xi'_x)$ is an ideal bundle of $h(\xi)$ where*

$$h_1(\xi'_x) = \{T \in h(\xi_x) \mid T(\xi_x) \subseteq \xi'_x\}$$

Proof. Each $h_1(\xi'_x)$ is an ideal in $h(\xi_x)$. It is enough to prove that $h_1(\xi')$ is a vector bundle. Since the bundle ξ is semisimple and ξ' is semisimple being its ideal bundle, ξ and ξ' are locally trivial by Theorem (3.4). So we have Jordan algebras J and J' and Jordan bundle isomorphisms $\Phi : U \times J \rightarrow \cup_{x \in X} \xi_x$ and $\Psi : U \times J' \rightarrow \cup_{x \in X} \xi'_x$. Then, $\Phi^{-1}\Psi : U \times J' \rightarrow U \times J$ is a vector bundle monomorphism satisfying the hypotheses of Lemma (4.1). Hence, there exists a finite open cover

$\{U_i\}$ for U such that $(\Phi^{-1}\Psi)(x, J) = (\Phi^{-1}\Psi)(x', J)$ for all $x, x' \in U_i$. Consequently, the neighbourhood U can be shrunk so that there exists an ideal I of J and such that Φ maps $U \times I$ onto $\cup_{x \in U} \xi'_x$. Let $h_1(I) = \{T \in h(I) \mid T(I) \subseteq I\}$. Given $T \in h_1(I)$, consider $g(\Phi)_x(T) = \Phi_x T \Phi_x^{-1}$.

$$\begin{aligned} g(\Phi)_x(T)(\xi_x) &= (\Phi_x T \Phi_x^{-1})(\xi_x) \\ &= (\Phi_x T)(J) \subseteq \Phi_x(I) = \xi'_x. \end{aligned}$$

Hence, $g(\Phi)_x(T) \in h_1(\xi'_x)$. That is, $g(\Phi)$ maps $U \times h_1(I)$ onto $\cup_{x \in U} h_1(\xi'_x)$. That is, $h_1(\xi')$ is an ideal sub-bundle of $h(\xi)$. \square

Theorem 4.3. *Every semisimple Jordan algebra bundle can be uniquely written as the direct sum of simple ideal bundles.*

Proof. Let ξ be a semisimple Jordan algebra bundle. Each fibre ξ_x has a unit element being a semisimple Jordan algebra [13, Theorem 4.7, p. 99]. Hence, the corresponding Lie algebra bundle $L(\xi)$ exists [1]. The semisimplicity of ξ_x implies that of $L(\xi_x)$ [7, p. 805] and so $L(\xi)$ is a semisimple Lie algebra bundle. Then, $L(\xi)$ can be written as follows:

$$L(\xi) = L_1 \oplus L_2 \oplus \dots \oplus L_n,$$

where each L_i is a simple ideal bundle of $L(\xi)$ [14]. Also each L_i is of the form $L_i = \cup_{x \in X} (h_i)_x \oplus (A_i)_x \oplus (B_i)_{x,}$ [7, Lemma 1, p. 789]. Each $\xi_i = \cup_{x \in X} (A_i)_x$ is an ideal bundle of ξ and each ξ_i is simple since L_i is simple. Hence, $\xi = \xi_1 \oplus \xi_2 \oplus \dots \oplus \xi_n$.

Let us prove the uniqueness of the decomposition. Let, ξ be expressed as follows:

$$\xi_1 \oplus \xi_2 \oplus \dots \oplus \xi_n = \xi = \xi'_1 \oplus \xi'_2 \oplus \dots \oplus \xi'_m,$$

where each ξ_i and ξ'_j are simple ideal bundles. Consider $\oplus_1^n h_1(\xi_i)_x$ where $h_1(\xi_i)_x = \{T \in h(\xi_x) \mid T(\xi_x) \subseteq (\xi_i)_x\}$. We have $\oplus_1^n h_1(\xi_i)_x \subseteq h(\xi_x)$. Let $L(a) \in h(\xi_x)$, then a can be written as $a = \oplus_1^n a_i$, $a_i \in (\xi_i)_x$. Since $(\xi_i)_x$ is an ideal in ξ_x , $L(a_i)$ maps ξ_x into $(\xi_i)_x$. So

$$L(a) = \oplus_{i=1}^n L(a_i) \in \oplus_1^n h_1(\xi_i)_x,$$

Let $[L(a), L(b)] \in h(\xi_x)$, where $a = \oplus_1^n a_i$ and $b = \oplus_1^n b_i$, $a_i, b_i \in (\xi_i)_x$. Since $[L(a_i), L(b_j)](u) \in (\xi_i)_x \cap (\xi_j)_x = 0$ for $i \neq j$, we obtain that

$$[L(a), L(b)] = \oplus_1^n [L(a_i), L(b_j)] \in \oplus_1^n h_1(\xi_i)_x.$$

Consequently, $h(\xi_x) \subseteq \oplus_1^n h_1(\xi_i)_x$. Therefore, $h(\xi) = h_1(\xi_1) \oplus h_1(\xi_2) \oplus \dots \oplus h_1(\xi_n)$. Similarly, $h(\xi) = \oplus_1^m h_1(\xi'_j)$. Hence

$$\begin{aligned} L(\xi) &= L_1(\xi_1) \oplus L_1(\xi_2) \oplus \dots \oplus L_1(\xi_n) \\ &= L_1(\xi'_1) \oplus L_1(\xi'_2) \oplus \dots \oplus L_1(\xi'_m) \end{aligned}$$

where $L_1(\xi_i) = h_1(\xi_i) \oplus \xi_i \oplus \xi_i$ and $L_1(\xi'_j) = h_1(\xi'_j) \oplus \xi'_j \oplus \xi'_j$ are ideal bundles of $L(\xi)$ by Lemma (4.2). Then by [14, Theorem 2.8], we obtain that $m = n$ and $L_1(\xi_i)$ coincides with one of the $L_1(\xi'_j)$. Then, obviously ξ'_j coincides with ξ_i except for the order. \square

References

1. Kumar R. On Wedderburn principal theorem for Jordan algebra bundles. Commun Algebra. 2021; 49(4): 1431-5.
2. Jordan P. Über eine Klasse nichtassoziativer hyperkomplexen Algebren. Gott Nachr; 1932. 569-75.

3. Jordan P. Uber Verallgemeinerungsmoglichkeiten des Formalismus der Quantenmechanik, Gott. Nachr; 1933. 209-17.
4. Jordan P. em Uber die Multiplikation quantenmechanischen Grossen. Z Phys. 1933; 80: 285-91.
5. McCrimmon K. Jordan algebras and their applications. Bull Amer Math Soc. 1978; 84: 612-27.
6. Prema G, Kiranagi BS. Lie algebras bundles defined by Jordan algebra bundles. Bull Math Soc Sci Math Roumanie. 1987; 31(79): 255-64.
7. Koecher M. Imbedding of Jordan algebras into Lie algebras-I. Amer J Math. 1967; 89: 787-816.
8. Atiyah MF. K-Theory. New York, Amsterdam: W.A.Benjamin, Inc.; 1967.
9. Finston DR. Rigidity and compact real forms of semisimple complex Jordan algebras. Commun Algebra. 1990; 18(10): 3323-38.
10. Arens R. Topologies for homeomorphism groups. Amer J Math. 1946; 68: 593-610.
11. Steenrod N. The topology of fibre bundles. Princeton, New Jersey: Princeton University Press; 1974.
12. Cohn PM. Lie Groups. Cambridge: Cambridge University Press; 1965.
13. Schafer RD. Introduction to non-associative algebras. New York: Acd. Press; 1966.
14. Kiranagi BS. Semisimple Lie algebra bundles. Bull Math Soc Sci Math Roumanie. 1983; 27(75): 253-7.

Further reading

15. Jacobson N. Structure and representations of Jordan algebras. Amer Math Soc Coll Publs. 1968.
16. Kiranagi BS, Prema G. A decomposition theorem of Lie algebra bundles. Comm Algebra. 1990; 18(6): 1869-77.
17. Koecher M. Imbedding of Jordan algebras into Lie algebras-II. Amer J Math. 1968; 90: 476-510.

Corresponding author

Ranjitha Kumar can be contacted at: ranju286math@gmail.com, ranjitha.kumar@reva.edu.in