

Biological properties in relation to health promotion effects of *Garcinia mangostana* (queen of fruit)

A short report

Naymul Karim and Jitbanjong Tangpong
*School of Allied Health Sciences, Walailak University,
Nakhon Si Thammarat, Thailand*

Abstract

Purpose – For the prevention and cure of disease, patient use various types of chemical and drug agents. Along with their curative effect, almost all drugs have some destructive effects and side-effects. Due to the minimal and/or none of unwanted side-effect, recently, the use of herbal remedy as the drug of choice becomes the preference choice. The mangosteen, *Garcinia mangostana*, contains various types of polyphenols. It has been used as a traditional medicine from the ancient times till present days. The purpose of this paper is to investigate the biological properties of mangosteen in relation to health promotion effects.

Design/methodology/approach – Several research papers from well-known database (such as PubMed, Google scholar, Scopus and Scencedirect) were reviewed without considering publication-times to understand the biological properties of mangosteen.

Findings – Mangosteen and its xanthone exerted diverse biological activities such as anti-oxidant, anti-inflammatory, anti-allergy, anti-bacteria, anti-fungal, anti-malaria, anticancer and anti-diabetes.

Originality/value – Based on these studies, mangosteen is beneficial dietary supplement of overall human health.

Keywords *Garcinia mangostana*, Xanthones, Biological properties

Paper type Short report

Introduction

The mangosteen, *Garcinia mangostana* (GM), which defines as “queen of fruits”[1], is native plant mainly found in the tropical rainforest areas of South-Asian countries, e.g., Myanmar, Thailand, Indonesia, Malaysia, Philippines, Sri-Lanka, etc. It belongs to the Guttiferace Family. GM has known and being used as traditional medicine to treat various types of diseases such as diarrhea, abdominal pain, dysentery, wound-infections and chronic ulcer[1]. According to various studies, GM extracts possess potent anti-oxidant[2], anti-tumoral[3], anti-allergic[4], anti-inflammatory[5] anti-bacterial[6] and anti-fungal activities[7]. Mangostana is famous for its tasty fruit named as mangosteen fruit. It is a dark purple/reddish color fruit with white soft and delicious consumable pulp. This pulp is little bit acidic and sweet in flavor with charming smell. Products containing mangosteen juice/extract are highly demandable in the beverage market; and in the USA, the sales of mangosteen products exceeded up to \$200m in 2008. A variety of secondary metabolites have been isolated from GM, xanthone is one of them[8]. Mangosteen fruit contains 160 aromatic compounds (epicarp) and 105 compounds (endocarp) evaluated by gas chromatography–mass spectral analysis[9].



Tang *et al.*[10], in their randomized, double-blind, placebo-controlled trial, reported that mangosteen intake for 30 days enhanced human immune responses compared to placebo controls. Udani *et al.*[11] found that intake of blended mangosteen juice by obese subjects for eight-weeks ameliorated the inflammation compared to placebo group. Another randomized, double-blind, placebo-controlled trial by Kudiganti *et al.*[12] revealed that consumption of meratrim (contained GM fruit rinds) for 16 weeks by healthy overweight subjects, significantly reduced the body weight, BMI, hip size and serum lipid profile compared to the placebo control. After ingestion of 60 mL mangosteen juice by healthy adult subjects with high-fat breakfast, xanthone were detected in serum approximately 762–4,030 nM/L/h and 0.9–11.1 μ M in 24 hours urine indicate the well-absorption of xanthone when taken with high-fat-diet[13]. Chang *et al.*[14] unveiled that acute administration of 250 mL mangosteen juice one hour before cycle ergometer exercise, does not have impact on exercise-mediated physical fatigue. The acute toxicity study of ethyl-acetate fraction shows no toxicity to the experimental rat evaluated by physical changes and mortality[15]. Therefore, GM and its isolated xanthones are safe for people.

Xanthone from GM

Xanthone (known as xanthen-9H-ones), an active compound, is an important element of oxygenated heterocycles group. It is the secondary metabolites obtained from higher plant family, fungi and lichen. It has been classified into five major groups: simple oxygenated xanthone, xanthone glycosides, prenylated xanthone, xanthone-lignoids and miscellaneous xanthone[8]. Maruganadan *et al.*[16] revealed that around 1,000 different xanthones have been isolated from the natural source. During the year of 2000 to 2004, about 278 new xanthones have been obtained from 20 higher plant families[17]. Mangostin (known as α -mangostin) is the first xanthone and was first isolated in 1855. Currently, 54 xanthones were isolated from the GM's pericarp, e.g., β -mangostin, γ -mangostin, mangostenol, 1-isomangostin, 1-isomangostin hydrate, etc[8].

Side effect of GM

There are few scientific reports published about mangosteen side-effect. Daily consumption of mangosteen juice for a long time may produce type-B lactate acidosis in chronic kidney disease patient and metabolic syndrome. This may be due to the releasing of cytochrome-c, by the α -mangostin, cytosol from the mitochondria and directly impair the mitochondrial electron transport that leads to lactic acidosis[18]. Liu *et al.*[19] found that α - and γ -mangostin from GM can inhibit platelet aggregation and induce cytolysis, therefore it should be avoided before any surgery.

Biological properties of GM

Anti-oxidant properties

Many scientists reported that GM and its isolated xanthone exhibited potent anti-oxidant properties[20–23]. In the cisplatin-induced nephrotoxic rats, the α -mangostin from GM exerted renoprotective effect by reducing cisplatin-induced renal oxidative/nitrosative stress[2]. Tjahjani *et al.*[20] evaluated the anti-oxidant activity of the different fraction of mangosteen rind, and found that the ethyl-acetate fraction exerted higher anti-oxidant than other fractions. The GM's pericarp extract also inhibited the formation of pentosidine and reduced advanced glycation end-products accumulation in the skin, which suppressed the glycation stress and improve skin conditions[21]. Xie *et al.*[22] found that drinking 245 mL mangosteen contained beverage by healthy populations enhanced the plasma anti-oxidant capacity (maximum 60 percent after one hour). A clinical study reported that oral administration of polar fraction extract from mangosteen pericarp to human subjects for 24-weeks enhanced the anti-oxidant activity without producing potential side-effect[23].

Anti-allergy and anti-inflammatory properties

Anti-allergy and anti-inflammatory properties of GM have been proved by various scientific evidence. The GM's pericarp inhibited the histamine- and serotonin-induced isolated thoracic rabbit aorta contractions, and blocked the histamine and serotonin receptor, respectively[4]. In RBL-2H3 cell line, 40 percent ethanol extract of mangosteen was found to inhibit the histamine release and prostaglandin E2 (PGE2) synthesis activities[24]. The 18 hours treatment of γ -mangostin suppressed the continuous PGE2 release and cyclooxygenase-2 (COX-2) gene expression in the C6 rat glioma cells in concentration-dependent manners[5]. The extracted isogarcinol from GM exhibited anti-inflammatory activity by suppressing the CD4 T-cells regulation in the murine model[25]. The combination of Garcinia with cocoa, coffee and green tea significantly reduced the lipid content from serum and liver in a dose-dependent manner. It also improved homeostasis model by assessing insulin resistance index and pro-inflammatory cytokines (TNF- α , IL-6)[26].

Anti-bacterial and anti-fungal properties

A number of studies described anti-bacterial and anti-fungal properties of xanthone extracts from GM[6–7, 27–30]. It was reported that the polysaccharides compound obtained from mangosteen fruit increased polymorphonuclear phagocytic cell activity against *Salmonella enteritidis*[27]. Prenylated xanthone from GM's pericarp has potential effect on tuberculosis, showed potent anti-bacterial activity by inhibiting *Mycobacterium tuberculosis*[28]. Sakagami *et al.*[6] found that α -mangostin inhibited the vancomycin-resistant *Enterococci* (VRE) and MRSA (MIC value 6.25 and 12.5 $\mu\text{g/ml}$). Different xanthenes (α -, γ -mangostin, gartanin, garcinone D, BR-xanthone and euxanthone) isolated from mangosteen exhibited the anti-fungal activity against phytopathogenic fungi (e.g. *Fusarium oxysporum vasinfectum*, *Alternaria tenuis*, and *Dreschlera oryzae*, etc.) by showing potent inhibitory activity[7]. Moreover, ethanol and chloroform extract of mangosteen demonstrated anti-bacterial activity, evaluated by colony formation and zone of inhibition of the *E. coli*, streptococcus and lactobacillus bacteria[29, 30].

Anti-malarial properties

The α -mangostin from GM exhibited *in-vitro* anti-malarial activity (IC_{50} values $17 \pm 1 \mu\text{M}$) against *Plasmodium falciparum* compared to chloroquine standard drug (IC_{50} values $0.59 \pm 0.02 \mu\text{M}$), evaluated on *P. falciparum*-infected erythrocytes[31]. Laphookhieo and his team isolated four xanthone (5-O-methylcelebixanthone, celebixanthone, cochinchinone C and β -mangostin) from the root of *Cratoxylum cochinchinense*, and found that β -mangostin exerted better *in-vitro* anti-malarial effect (IC_{50} value 7.2) than other compound (IC_{50} values 3.2, 4.9 and 2.6 mg/ml, respectively), when they were evaluated via [^3H]-hypoxanthine uptake by *P. falciparum*[32]. An *in-vivo* study exposed that α -mangostin ($\text{IC}_{50} = 0.2 \pm 0.01 \mu\text{M}$) showed more effective anti-malarial activity than γ -mangostin ($\text{IC}_{50} = 121.2 \pm 1.0 \mu\text{M}$) against chloroquine resistant strain of *P. falciparum* on the malarial murine model[33]. In addition, different fraction of mangosteen exerted synergistic anti-malarial activity with artemisinin drug against *Plasmodium falciparum* 3D7 clone[34].

Anti-tumoral properties

Several studies revealed the anticancer activities of xanthenes derivative from GM[35–41]. In HL60 cell line, six xanthenes (α -, β -, γ -mangostin, mangostinone, garcinone E and 2-isoprenyl-1, 7-dihydroxy-3-methoxy xanthone) possessed the cytotoxic effect by inhibiting cell growth, while the α -mangostin exerted the highest inhibitory activity than others[3]. The α -mangostin also induced the mitochondrial dysfunction in the early phase through activating caspases-3/9, ROS production, and cytochrome-c release[35]. Nakagawa *et al.*[36]

unveiled that the α -mangostin exerted *in-vitro* cytotoxic effect against DLD-1 cell line. In HeLa cell line, encapsulation of GM with methyl-cellulose and ethyl-cellulose nanoparticles showed two-fold higher anticancer activity than the encapsulation of GM with ethyl-cellulose[37]. The α -mangostin exhibited the most potent mammalian DNA polymerase inhibition, evaluated by the inhibition of human DNA topoisomerases I and II activities (IC_{50} values 15.0 and $7.5 \mu M$). It also suppressed the HCT116 cell proliferation (LC_{50} value $18.5 \mu M$)[38]. In prostate cancer, gartanin xanthone bind with androgen receptors (AR) and enhance AR degradation[39]. Manimekalai *et al.*[40] revealed that mangosteen showed anticancer activity on HepG2 cell line evaluated by the cell viability assay. A total of 14 isolated compounds from chloroform fraction of mangosteen exerted anticancer activity against different cancer cell line, e.g., HepG2, HCT116 and MCF7[41].

Anti-diabetic properties

Different fractions of the GM's pericarp and isolated xanthone showed an anti-diabetic effect by inhibiting α -amylase and α -glucosidase enzyme activities[42, 43]. Ethanolic extract of GM showed the anti-diabetic effect against streptozotocin-induced diabetic rats by lowering high blood glucose, biochemical parameters, hepatic architectures and increased the HDL and total protein levels, compared to the control group[44]. While, in high-fat-diet- and streptozotocin-induced diabetic mice, mangosteen vinegar rind from GM, lowered the hyperglycemia, hyperlipidemia, oxidative stress marker, hepatic damage marker and improved the glycogen contents and anti-oxidant markers[45]. In similar model, isolated xanthone from GM demonstrated the nephroprotective effect by reducing the body weight, high glucose level, kidney hypertrophy, kidney damage marker and MDA level of plasma and kidney tissue[46].

Conclusion

Recently, the increasing awareness on the negative impacts of nutrition and diets, such as causing obesity, development of chronic diseases, cardiovascular disease, diabetes, cancer and other disease, was quite alarming as well as obvious especially in the western countries. A large number of bioactive compounds have already been identified in foods and drinks, contain high percentage of polyphenolic compounds (e.g. xanthone, lignans, phenolic acids, etc.).

The GM possesses potential biological properties without producing noticeable side-effect. It boosts up immune systems, reduced metabolic syndromes and its associated diseases, respectively. The GM's xanthone was found to associate with health beneficial effects by lowering oxidative stress, reducing inflammation, managing and controlling obesity and diabetes, inhibiting the growth of cancer cells as well as showing anti-allergic, anti-bacterial, anti-fungal and anti-malarial properties.

However, most of investigations on GM mentioned in this report were conducted either in cell-free, cell-type or animal model and none of them was on humans. Therefore, the further studies of health effects on human subjects are crucial and needed.

References

1. Cui J, Hu W, Cai Z, Liu Y, Li S, Tao W, *et al.* New medicinal properties of mangostins: analgesic activity and pharmacological characterization of active ingredients from the fruit hull of *Garcinia mangostana* L. *Pharmacol Biochem Behav.* 2010 Apr; 95(2): 166-72. doi: 10.1016/j.pbb.2009.12.021
2. Perez-Rojas JM, Cruz C, Garcia-Lopez P, Sanchez-Gonzalez DJ, Martinez-Martinez CM, Ceballos G, *et al.* Renoprotection by alpha-Mangostin is related to the attenuation in renal oxidative/nitrosative stress induced by cisplatin nephrotoxicity. *Free Radic Res.* 2009; 43(11): 1122-32.

3. Matsumoto K, Akao Y, Kobayashi E, Ohguchi K, Ito T, Tanaka T, *et al.* Induction of apoptosis by xanthenes from mangosteen in human leukemia cell lines. *J Nat Prod.* 2003 Aug; 66(8): 1124-7. doi: 10.1021/np020546u
4. Chairungsrilerd N, Furukawa K, Ohta T, Nozoe S, Ohizumi Y. Histaminergic and serotonergic receptor blocking substances from the medicinal plant *Garcinia mangostana*. *Planta Med.* 1996 Oct; 62(5): 471-2. doi: 10.1055/s-2006-957943
5. Nakatani K, Yamakuni T, Kondo N, Arakawa T, Oosawa K, Shimura S, *et al.* γ -Mangostin inhibits inhibitor- κ B kinase activity and decreases lipopolysaccharide-induced cyclooxygenase-2 gene expression in C6 rat glioma cells. *Mol Pharmacol.* 2004 Sep; 66(3): 667-74. doi: 10.1124/mol.104.002626
6. Sakagami Y, Iinuma M, Piyasena KG, Dharmaratne HR. Antibacterial activity of α -mangostin against Vancomycin Resistant Enterococci (VRE) and synergism with antibiotics. *Phytomedicine.* 2005 Mar; 12(3): 203-8.
7. Gopalakrishnan G, Banumathi B, Suresh G. Evaluation of the antifungal activity of natural xanthenes from *Garcinia mangostana* and their synthetic derivatives. *J Nat Prod.* 1997 May; 60(5): 519-24. doi: 10.1021/np970165u
8. Gutierrez-Orozco F, Failla ML. Biological activities and bioavailability of mangosteen xanthenes: a critical review of the current evidence. *Nutrients.* 2013 Aug; 5(8): 3163-83. doi: 10.3390/nu5083163
9. Manimekalai I, Sivakumari K, Ashok K, Rajesh S. Phytochemical profiling of mangosteen fruit, *Garcinia mangostana*. *World J Pharm Pharm Sci.* 2016; 5(2): 221-52.
10. Tang YP, Li PG, Kondo M, Ji HP, Kou Y, Ou B Effect of a mangosteen dietary supplement on human immune function: a randomized, double-blind, placebo-controlled trial. *J Med Food.* 2009 Aug; 12(4): 755-63. doi: 10.1089/jmf.2008.0204
11. Udani JK, Singh BB, Barrett ML, Singh VJ. Evaluation of mangosteen juice blend on biomarkers of inflammation in obese subjects: a pilot, dose finding study. *Nutr J.* 2009 Oct; 8(1): 48. doi: 10.1186/1475-2891-8-48
12. Kudiganti V, Kodur RR, Kodur SR, Halemane M, Deep DK. Efficacy and tolerability of Meratrim for weight management: a randomized, double-blind, placebo-controlled study in healthy overweight human subjects. *Lipids Health Dis.* 2016 Aug; 15(1): 136. doi: 10.1186/s12944-016-0306-4
13. Chitchumroonchokchai C, Riedl KM, Suksumrarn S, Clinton SK, Kinghorn AD, Failla ML. Xanthenes in mangosteen juice are absorbed and partially conjugated by healthy adults. *J Nutr.* 2012 Apr; 142(4): 675-80. doi: 10.3945/jn.111.156992
14. Chang CW, Huang TZ, Chang WH, Tseng YC, Wu YT, Hsu MC. Acute *Garcinia mangostana* (mangosteen) supplementation does not alleviate physical fatigue during exercise: a randomized, double-blind, placebo-controlled, crossover trial. *J Int Soc Sports Nutr.* 2016; 13(1): 20. doi: 10.1186/s12970-016-0132-0
15. Rahmayanti F, Suniarti DF, Masud ZA, Bachtiar BM, Wimardhani YS, Subita GP. Acute oral toxicity testing of ethyl acetate fraction from *Garcinia mangostana* Linn extract in sprague-dawley rats. *Res J Med Plants.* 2016; 10(3): 261-4. doi: 10.3923/rjmp.2016.261.264
16. Muruganandan S, Srinivasan K, Gupta S, Gupta PK, Lal J. Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. *J Ethnopharmacol.* 2005 Mar; 97(3): 497-501. doi: 10.1016/j.jep.2004.12.010
17. Sousa ME, Pinto MM. Synthesis of xanthenes: an overview. *Curr Med Chem.* 2005; 12(21): 2447-79.
18. Wong LP, Klemmer PJ. Severe lactic acidosis associated with juice of the mangosteen fruit *Garcinia mangostana*. *Am J Kidney Dis.* 2008 May; 51(5): 829-33. doi: 10.1053/j.ajkd.2007.12.043
19. Liu Y, Park JM, Chang KH, Chin YW, Lee MY. α - and γ -mangostin cause shape changes, inhibit aggregation and induce cytolysis of rat platelets. *Chem Biol Interact.* 2015 Oct; 240: 240-8. doi: 10.1016/j.cbi.2015.08.021
20. Tjahjani S, Widowati W, Khiong K, Suhendra A, Tjokropranoto R. Antioxidant properties of *Garcinia mangostana* L (mangosteen) rind. *Procedia Chem.* 2014; 13: 198-203. doi: 10.1016/j.proche.2014.12.027

21. Ohno R, Moroishi N, Sugawa H, Maejima K, Saigusa M, Yamanaka M, et al. Mangosteen pericarp extract inhibits the formation of pentosidine and ameliorates skin elasticity. *J Clin Biochem Nutr.* 2015 Jul; 57(1): 27-32. doi: 10.3164/jcbn.15-13
22. Xie Z, Sintara M, Chang T, Ou B. Functional beverage of *Garcinia mangostana* (mangosteen) enhances plasma antioxidant capacity in healthy adults. *Food Sci Nutr.* 2015 Jan; 3(1): 32-8. doi: 10.1002/fsn3.187
23. Suthammarak W, Numpraphrut P, Charoensakdi R, Neungton N, Tunrungruangtavee V, Jaisupa N, et al. Antioxidant-enhancing property of the polar fraction of mangosteen pericarp extract and evaluation of its safety in humans. *Oxid Med Cell Longev.* 2016 Jul; 2016: 12511. doi: 10.1155/2016/1293036
24. Nakatani K, Atsumi M, Arakawa T, Oosawa K, Shimura S, Nakahata N, et al. Inhibitions of histamine release and prostaglandin E2 synthesis by mangosteen, a Thai medicinal plant. *Biol Pharm Bull.* 2002 Sep; 25(9): 1137-41.
25. Li W, Li H, Zhang M, Zhong Y, Wang M, Cen J, et al. Isogarcinol extracted from *Garcinia mangostana* L. ameliorates systemic lupus erythematosus-like disease in a murine model. *J Agric Food Chem.* 2015 Sep; 63(38): 8452-9. doi: 10.1021/acs.jafc.5b03425
26. Chang CW, Hsu YJ, Chen YM, Huang WC, Huang CC, Hsu MC. Effects of combined extract of cocoa, coffee, green tea and garcinia on lipid profiles, glycaemic markers and inflammatory responses in hamsters. *BMC Complement Altern Med.* 2015 Aug; 15(1): 269. doi: 10.1186/s12906-015-0806-1
27. Chanarat P, Chanarat N, Fujihara M, Nagumo T. Immunopharmacological activity of polysaccharide from the pericarp of mangosteen garcinia: phagocytic intracellular killing activities. *J Med Assoc Thai.* 1997 Sep; 80(S1): S149-54.
28. Suksamrarn S, Suwannapoch N, Phakhodee W, Thanuhiranlert J, Ratananukul P, Chimnoi N, et al. Antimycobacterial activity of prenylated xanthenes from the fruits of *Garcinia mangostana*. *Chem Pharm Bull (Tokyo).* 2003 Jul; 51(7): 857-9.
29. Soetikno JS, Handayani R, Rahayu RD. Assay for antimicrobial activity of mangosteen rind extracts. *Journal of Innovations in Pharmaceuticals and Biological Sciences.* 2016; 3(1): 54-60.
30. Janardhanan S, Mahendra J, Giriya AS, Mahendra L, Priyadharsini V. Antimicrobial effects of *Garcinia mangostana* on cariogenic microorganisms. *J Clin Diagn Res.* 2017 Jan; 11(1): 19-22. doi: 10.7860/jcdr/2017/22143.9160
31. Mahabusarakam W, Kuaha K, Wilairat P, Taylor WC. Prenylated xanthenes as potential antiplasmodial substances. *Planta Med.* 2006 Aug; 72(10): 912-6. doi: 10.1055/s-2006-947190
32. Laphookhieo S, Syers JK, Kiattansakul R, Chantrapromma K. Cytotoxic and antimalarial prenylated xanthenes from *Cratoxylum cochinchinense*. *Chem Pharm Bull (Tokyo).* 2006 May; 54(5): 745-7.
33. Upegui Y, Robledo SM, Gil Romero JF, Quinones W, Archbold R, Torres F, et al. *In vivo* antimalarial activity of α -mangostin and the new xanthone δ -mangostin. *Phytother Res.* 2015 Aug; 29(8): 1195-201. doi: 10.1002/ptr.5362
34. Tjahjani S. Antimalarial activity of *Garcinia mangostana* L rind and its synergistic effect with artemisinin *in vitro*. *BMC Complement Altern Med.* 2017 Feb; 17(1): 131. doi: 10.1186/s12906-017-1649-8
35. Matsumoto K, Akao Y, Yi H, Ohguchi K, Ito T, Tanaka T, et al. Preferential target is mitochondria in alpha-mangostin-induced apoptosis in human leukemia HL60 cells. *Bioorg Med Chem.* 2004 Nov; 12(22): 5799-806. doi: 10.1016/j.bmc.2004.08.034
36. Nakagawa Y, Inuma M, Naoe T, Nozawa Y, Akao Y. Characterized mechanism of α -mangostin-induced cell death: caspase-independent apoptosis with release of endonuclease-G from mitochondria and increased miR-143 expression in human colorectal cancer DLD-1 cells. *Bioorg Med Chem.* 2007 Aug 15; 15(16): 5620-8. doi: 10.1016/j.bmc.2007.04.071
37. Pan-In P, Wanichwecharungruang S, Hanes J, Kim AJ. Cellular trafficking and anticancer activity of *Garcinia mangostana* extract-encapsulated polymeric nanoparticles. *Int J Nanomedicine.* 2014; 9: 3677-86. doi: 10.2147/ijn.S66511

38. Mizushina Y, Kuriyama I, Nakahara T, Kawashima Y, Yoshida H. Inhibitory effects of α -mangostin on mammalian DNA polymerase, topoisomerase, and human cancer cell proliferation. *Food Chem Toxicol.* 2013 Sep; 59: 793-800. doi: 10.1016/j.fct.2013.06.027
39. Li G, Petiwala SM, Yan M, Won JH, Petukhov PA, Johnson JJ. Gartanin, an isoprenylated xanthone from the mangosteen fruit (*Garcinia mangostana*), is an androgen receptor degradation enhancer. *Mol Nutr Food Res.* 2016 Jun; 60(6): 1458-69. doi: 10.1002/mnfr.201600037
40. Manimekalai I, Sivakumari K, Ashok K, Rajesh S. Antioxidant and anticancer potential of mangosteen fruit, *Garcinia mangostana* against hepatocellular carcinoma (HEPG-2) cell line. *World J Pharm Pharm Sci.* 2016; 5(2): 253-93.
41. Mohamed GA, Al-Abd AM, El-Halawany AM, Abdallah HM, Ibrahim SRM. New xanthones and cytotoxic constituents from *Garcinia mangostana* fruit hulls against human hepatocellular, breast, and colorectal cancer cell lines. *J Ethnopharmacol.* 2017 Feb; 198: 302-12. doi: 10.1016/j.jep.2017.01.030
42. Loo AE, Huang D. Assay-guided fractionation study of α -amylase inhibitors from *Garcinia mangostana* pericarp. *J Agric Food Chem.* 2007 Nov; 55(24): 9805-10. doi: 10.1021/jf071500f
43. Ryu HW, Cho JK, Curtis-Long MJ, Yuk HJ, Kim YS, Jung S, et al. α -Glucosidase inhibition and antihyperglycemic activity of prenylated xanthones from *Garcinia mangostana*. *Phytochemistry.* 2011 Dec; 72(17): 2148-54. doi: 10.1016/j.phytochem.2011.08.007
44. Taher M, Tg Zakaria TM, Susanti D, Zakaria ZA. Hypoglycaemic activity of ethanolic extract of *Garcinia mangostana* Linn. in normoglycaemic and streptozotocin-induced diabetic rats. *BMC Complement Altern Med.* 2016 May; 16(1): 135. doi: 10.1186/s12906-016-1118-9
45. Karim N, Jeenduang N, Tangpong J. Anti-glycemic and anti-hepatotoxic effects of mangosteen vinegar rind from *Garcinia mangostana* against HFD/STZ-induced type II diabetes in mice. *Pol J Food Nutr Sci.* 2018; 68(2): 163-9. doi: 10.1515/pjfn-2017-0018
46. Karim N, Jeenduang N, Tangpong J. Renoprotective Effects of xanthone derivatives from *Garcinia mangostana* against high fat diet and streptozotocin-induced type II diabetes in mice. *Walailak J Sci Technol.* 2018; 15(2): 107-16.

Corresponding author

Jitbanjong Tangpong can be contacted at: njibjoy@yahoo.com